



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>XXXX XXX/XXXXX</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/47284</b> <b>(43) International Publication Date:</b> 17 August 2000 (17.08.2000)
<b>(21) International Application Number:</b> PCT/EP00/01147 <b>(22) International Filing Date:</b> 10 February 2000 (10.02.2000) <b>(30) Priority Data:</b> 9903265.8 12 February 1999 (12.02.1999) GB 9903278.1 12 February 1999 (12.02.1999) GB 9903282.3 12 February 1999 (12.02.1999) GB 9903284.9 12 February 1999 (12.02.1999) GB 9906061.8 17 March 1999 (17.03.1999) GB <b>(60) Parent Application or Grant</b> SMITHKLINE BEECHAM P.L.C. [/]; (). IRVING, Elaine, Alison [/]; (). SANGER, Gareth, John [/]; (). IRVING, Elaine, Alison [/]; (). SANGER, Gareth, John [/]; (). BLAKEY, Alison, Jane ; ().		<b>Published</b>
<b>(54) Title: NOVEL USE OF OREXIN RECEPTOR ANTAGONISTS</b> <b>(54) Titre: NOUVELLE UTILISATION DES ANTAGONISTES DES RECEPTEURS DE L'OREXINE</b>  <b>(57) Abstract</b> <p>The use of orexin receptor antagonists as neuroprotectants, and in the treatment of nausea and vomiting, irritable bowel syndrome and other conditions associated with visceral pain.</p> <b>(57) Abrégé</b> <p>L'invention concerne l'utilisation des antagonistes des récepteurs de l'orexine comme neuroprotecteurs, ainsi que dans le cadre du traitement des nausées et vomissements, des colopathies fonctionnelles et autres pathologies associées à une douleur viscérale.</p>		

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : <b>A61P</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/47284</b> (43) International Publication Date: 17 August 2000 (17.08.00)
(21) International Application Number: <b>PCT/EP00/01147</b> (22) International Filing Date: 10 February 2000 (10.02.00)		(74) Agent: BLAKEY, Allison, Jane; SmithKline Beecham Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).	
(30) Priority Data: 9903282.3 12 February 1999 (12.02.99) GB 9903278.1 12 February 1999 (12.02.99) GB 9903284.9 12 February 1999 (12.02.99) GB 9903265.8 12 February 1999 (12.02.99) GB 9906061.8 17 March 1999 (17.03.99) GB		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): IRVING, Elaine, Alison [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). SANGER, Gareth, John [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).		<b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: NOVEL USE OF OREXIN RECEPTOR ANTAGONISTS			
(57) Abstract <p>The use of orexin receptor antagonists as neuroprotectants, and in the treatment of nausea and vomiting, irritable bowel syndrome and other conditions associated with visceral pain.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## Description

5

10

15

20

25

30

35

40

45

50

55

# NOVEL USE OF OREXIN RECEPTOR ANTAGONISTS

The present invention relates to the use of orexin receptor antagonists as neuroprotectants, and in the treatment of nausea and vomiting, irritable bowel syndrome and other conditions associated with visceral pain.

Many medically significant biological processes are mediated by proteins participating in signal transduction pathways that involve G-proteins and/or second messengers.

Polypeptides and polynucleotides encoding the human 7-transmembrane G-protein coupled neuropeptide receptor, orexin-1 (HFGAN72), have been identified and are disclosed in EP-A-875565, EP-A-875566 and WO 96/34877. Polypeptides and polynucleotides encoding a second human orexin receptor, orexin-2 (HFGANP), have been identified and are disclosed in EP-A-893498.

Polypeptides and polynucleotides encoding polypeptides which are ligands for the orexin-1 receptor, e.g. orexin-A (Lig72A) are disclosed in EP-A-849361.

Orexin receptors are found in the mammalian host and may be responsible for many biological functions.

Experiments have shown that central administration of the ligand orexin-A (described in more detail below) stimulated food intake in freely-feeding rats during a 4 hour time period. This increase was approximately four-fold over control rats receiving vehicle. These data suggest that orexin-A may be an endogenous regulator of appetite. Therefore, antagonists of its receptor may be useful in the treatment of obesity and diabetes, see *Cell*, 1998, 92, 573-585.

Rat sleep/EEG studies have also shown that central administration of orexin-A, an agonist of the orexin receptors, causes a dose-related increase in arousal, largely at the expense of a reduction in paradoxical sleep and slow wave sleep 2, when administered at the onset of the normal sleep period. Therefore antagonists of its receptor may be useful in the treatment of sleep disorders including insomnia.

Orexin-1 receptor mRNA and immunoreactivity is widely distributed throughout the brain including regions such as the hypothalamus, hippocampus and various other limbic and cortical areas. Thus the receptor is found in brain areas which have been reported to be affected by stroke in man and experimental stroke animals and head injury in man and experimental animals, to be involved in emesis and to be involved in visceral pain.

International Patent Applications PCT/GB98/02437 and PCT/EP99/03100 disclose various phenyl urea derivatives as orexin receptor antagonists.

It has now been discovered that compounds possessing orexin receptor antagonist activity, in particular orexin-1 receptor antagonist activity, also demonstrate neuroprotectant properties. Therefore, such compounds are useful in the treatment and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events in mammals, such as humans, including cerebral ischaemia after cardiac arrest, stroke and multi-infarct dementia, cerebral ischaemic events such as those resulting from surgery and/or during childbirth, and disorders resulting from traumatic head injury and subsequent haemorrhaging.

In particular the compounds are useful in the treatment and/or prophylaxis of stroke (ischaemic or haemorrhagic) or sub-arachnoid haemorrhage e.g. sub-arachnoid haemorrhage associated with traumatic head injury.

5 It has also been discovered that compounds possessing orexin receptor antagonist activity, in particular orexin-1 receptor antagonist activity, are useful for blocking the emetic response. The compounds are therefore of use in the treatment of nausea and vomiting associated with cancer therapy. The compounds are also of potential use in the treatment of post-operative or motion sickness.

10 It has also been discovered that compounds possessing orexin receptor antagonist activity, in particular orexin-1 receptor antagonist activity, may inhibit pain via an action on the peripheral C-fibres and are therefore useful in the treatment of conditions associated with visceral pain, such as irritable bowel syndrome and also migraine, angina and urge type incontinence such as inappropriate urge type incontinence associated with cystitis.

15 The present invention provides the use of an orexin receptor antagonist in the manufacture of a medicament for the treatment and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events, the treatment and/or prophylaxis of nausea and vomiting, or the treatment and/or prophylaxis of irritable bowel syndrome or other conditions associated with visceral pain.

20 The present invention also provides a method for the treatment and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events, the treatment and/or prophylaxis of nausea and vomiting, or the treatment and/or prophylaxis of irritable bowel syndrome or other conditions associated with visceral pain in mammals which comprises administering to a host in need thereof an effective amount of an orexin receptor antagonist or a pharmaceutically acceptable salt thereof.

25 When used in therapy, the orexin receptor antagonists are usually formulated in a standard pharmaceutical composition. Such compositions can be prepared using standard procedures.

30 The orexin receptor antagonist is preferably an orexin-1 receptor antagonist.

Orexin receptor antagonists may be administered by any convenient method, for example by oral, parenteral, buccal, sublingual, nasal, rectal or transdermal administration and the pharmaceutical compositions adapted accordingly.

35 Orexin receptor antagonists which are active when given orally can be formulated as liquids or solids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

40 A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable liquid carrier(s) for example an aqueous solvent such as water, ethanol or glycerine, or a non-aqueous solvent, such as polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring and/or colouring agent.

45 A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

50 A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

Compositions for nasal administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a propellant which can be a compressed gas such as compressed air or an organic propellant such as a fluorochlorohydrocarbon or hydrofluorocarbon. The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal administration are conveniently in the form of suppositories containing a conventional suppository base such as cocoa butter.

Compositions suitable for transdermal administration include ointments, gels and patches.

Preferably the composition is in unit dose form such as a tablet, capsule or ampoule.

The dose of the orexin receptor antagonist used in the treatment or prophylaxis of the abovementioned disorders or diseases will vary in the usual way with the particular disorder or disease being treated, the weight of the subject and other similar factors. However as a general rule suitable unit doses may be 0.05 to 1000 mg, more suitably 0.05 to 500 mg; such unit doses may be administered more than once a day for example two or three times a day, so that the total daily dosage is in the range of about 0.01 to 100 mg/kg; and such therapy may extend for a number of weeks or months.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

#### Experimental methods

Human orexin-A, referred to above, has the amino acid sequence:

pyroGlu	Pro	Leu	Pro	Asp	Cys	Cys	Arg	Gln	Lys	Thr	Cys	Ser	Cys	Arg	Leu
1			5				10				15				
Tyr	Glu	Leu	Leu	His	Gly	Ala	Gly	Asn	His	Ala	Ala	Gly	Ile	Leu	Thr
			20				25					30			
Leu-NH <sub>2</sub>															

Orexin-A can be employed in a process for screening for compounds (antagonists) which inhibit the ligand's activation of the orexin-1 receptor.

In general, such screening procedures involve providing appropriate cells which express the orexin-1 receptor on the surface thereof. Such cells include cells from mammals, yeast, *Drosophila* or *E. coli*. In particular, a polynucleotide encoding the orexin-1 receptor is employed to transfect cells to thereby express the receptor. The expressed receptor is then contacted with a test compound and an orexin-1 receptor ligand to observe inhibition of a functional response.

One such screening procedure involves the use of melanophores which are transfected to express the orexin-1 receptor. Such a screening technique is described in WO 92/01810.

Another such screening technique involves introducing RNA encoding the orexin-1 receptor into *Xenopus* oocytes to transiently express the receptor. The receptor oocytes may then be contacted with a receptor ligand and a compound to be screened, followed by detection of inhibition of a signal in the case of screening for compounds which are thought to inhibit activation of the receptor by the ligand.

Another method involves screening for compounds which inhibit activation of the receptor by determining inhibition of binding of a labelled orexin-1 receptor ligand to cells which have the receptor on the surface thereof. Such a method involves transfecting a eukaryotic cell with DNA encoding the orexin-1 receptor such that the cell expresses the receptor on its surface and contacting the cell or cell membrane preparation with a compound in the presence of a labelled form of an orexin-1 receptor ligand. The ligand can be labelled, e.g. by radioactivity. The amount of labelled ligand bound to the receptors is measured, e.g. by measuring radioactivity of the receptors. If the compound binds to the receptor as determined by a reduction of labelled ligand which binds to the receptors, the binding of labelled ligand to the receptor is inhibited.

Yet another screening technique involves the use of FLIPR equipment for high throughput screening of test compounds that inhibit mobilisation of intracellular calcium ions, or other ions, by affecting the interaction of an orexin-1 receptor ligand with the orexin-1 receptor. The ligand used in the screening method described below to determine orexin-1 receptor antagonist activity is orexin-A which has the amino acid sequence shown above.

#### Determination of Orexin-1 Receptor Antagonist Activity

The orexin-1 receptor antagonist activity of the compounds of formula (I) was determined in accordance with the following experimental method.

HEK293 cells expressing the human orexin-1 receptor were grown in cell medium (MEM medium with Earl's salts) containing 2 mM L-Glutamine, 0.4 mg/mL G418 Sulfate from GIBCO BRL and 10% heat inactivated fetal calf serum from Gibco BRL. The cells were seeded at 20,000 cells/100  $\mu$ L/well into 96-well black clear bottom sterile plates from Costar which had been pre-coated with 10  $\mu$ g/well of poly-L-lysine from SIGMA. The seeded plates were incubated overnight at 37°C in 5% CO<sub>2</sub>.

Agonists were prepared as 1 mM stocks in water:DMSO (1:1). EC<sub>50</sub> values (the concentration required to produce 50% maximal response) were estimated using 11x half log unit dilutions (Biomek 2000, Beckman) in Tyrode's buffer containing probenecid (10 mM HEPES with 145mM NaCl, 10mM glucose, 2.5 mM KCl, 1.5 mM CaCl<sub>2</sub>, 1.2 mM MgCl<sub>2</sub> and 2.5mM probenecid; pH7.4). Antagonists were prepared as 10 mM stocks in DMSO (100%). Antagonist IC<sub>50</sub> values (the concentration of compound needed to inhibit 50% of the agonist response) were



determined against 3.0 nM human orexin-A using 11x half log unit dilutions in Tyrode's buffer containing 10% DMSO and probenecid.

On the day of assay 50 µl of cell medium containing probenecid (Sigma) and Fluo3AM (Texas Fluorescence Laboratories) was added (Quadra, Tomtec) to each well to give final concentrations of 2.5 mM and 4 µM, respectively. The 96-well plates were incubated for 90 min at 37°C in 5% CO<sub>2</sub>. The loading solution containing dye was then aspirated and cells were washed with 4x 150 µl Tyrode's buffer containing probenecid and 0.1% gelatin (Denley Cell Wash). The volume of buffer left in each well was 125 µl. Antagonist or buffer (25 µl) was added (Quadra) the cell plates gently shaken and incubated at 37°C in 5% CO<sub>2</sub> for 30 min. Cell plates were then transferred to the Fluorescent Imaging Plate Reader (FLIPR, Molecular Devices) instrument and maintained at 37°C in humidified air. Prior to drug addition a single image of the cell plate was taken (signal test), to evaluate dye loading consistency. The run protocol used 60 images taken at 1 second intervals followed by a further 24 images at 5 second intervals. Agonists were added (by the FLIPR) after 20 sec (during continuous reading). From each well, peak fluorescence was determined over the whole assay period and the mean of readings 1-19 inclusive was subtracted from this figure. The peak increase in fluorescence was plotted against compound concentration and iteratively curve fitted using a four parameter logistic fit (as described by Bowen and Jerman, 1995, *TIPS*, 16, 413-417) to generate a concentration effect value. Antagonist Kb values were calculated using the equation:

$$K_b = IC_{50} / (1 + (3/EC_{50}))$$

where EC<sub>50</sub> was the potency of human orexin-A determined in the assay (in nM terms) and IC<sub>50</sub> is expressed in molar terms.

#### Pharmacological Models

##### Stroke and/or Head Injury

The following method may be used to demonstrate the neuroprotectant efficacy of orexin receptor antagonists in a model of stroke and/or head injury.

Male rats are anaesthetised and focal ischaemia induced via permanent middle cerebral artery (MCA) occlusion by a method as described by Longa *et al* (1989) *Stroke*, 20, 84-91. In a model of head injury the ischaemic effects of traumatic head injury may also be simulated by causing trauma to the head. Twenty-four hours after MCA occlusion, rats are perfusion fixed with neutral buffered formalin (NBF) containing 5% sucrose. The brains are then serially sectioned (1.5mm intervals) through the cerebrum from the anterior poles to the cerebellum and the sections (50 µm) stained with Cresyl Fast Violet. The sections are examined and the areas of ischaemic damage delineated at 8 pre-selected coronal levels from anterior + 3.0mm to posterior -7.5mm relative to bregma. Volumes are then corrected for hemispheric swelling. The test compound or vehicle is administered intravenously one hour post-MCA occlusion.

##### Inhibition of Chemotherapy Induced Nausea and Vomiting

Adult male ferrets (initial body weight 1.3-1.8kg) were individually housed, fed once daily (200 g Chum Puppy Food plus 50 g Lab Diet A) and were supplied with water *ad libitum*. Additionally, each animal was given ¼ pint of milk daily during an acclimatization period.

To facilitate intravenous administration of drugs, a chronic indwelling catheter was surgically implanted into the jugular vein using a modification of the technique described by Florczyk and Schurig, 1981 (*Pharmacol. Biochem. Behav.*, **14**, 255-257). Prior to surgery each animal was sedated with ketamine hydrochloride (40mg/animal intramuscularly) and anaesthetised with a halothane-N<sub>2</sub>O-O<sub>2</sub> mixture. A four day recovery period was allowed before commencement of an experiment.

For each group of animals, a preliminary study was carried out to establish an intravenous dose level of cytotoxic agent(s) which would give a consistent and reproducible emetic response. The appropriate dose of example was administered intravenously twice to each animal; 30 minutes before and 30-45 minutes after dosing with cytotoxic agent(s). Running controls received vehicle and cytotoxic agent(s) only.

A single emetic response commenced when an animal assumed a characteristic posture with retching and was concluded when either vomitus was expelled or was present in the mouth as evidenced by a chewing movement. The total number of emetic responses was determined during the four hour period following administration of cytotoxic agent(s) and the number of animals completely protected from emesis determined for each treatment.

#### Urinary Bladder Incontinence

Micturition reflexes are measured in animals in response to intraluminal bladder distension. There are several methods by which the sensitivity to normal levels of bladder distension are increased, thereby inducing bladder hyperreactivity and hence, providing a means by which the beneficial effect of orexin-1 receptor antagonists can be measured. Usually, these involve the application of a particular pathology to a rat or other species. In one example, the frequency of bladder voiding reflexes are increased in anaesthetized rats by chemical irritation of the bladder, using agents such as acetic acid or capsaicin (e.g. Yu and de Groat, *Brain Research*, 1998, **807**, 11-18); voiding reflexes are measured by recording intravesical pressure after catheterization of the bladder. Inhibitors of orexin-1 receptor-mediated function prevent this increase in bladder sensitivity.

#### Visceral Pain

The potential anti-migraine utility of orexin receptor antagonists may be evaluated in a range of models thought to represent the development of migraine (Chen *et al.*, 1999 *Bioorg. Med. Chem. Lett.*, 1999, **9**: 285-290; Raval *et al.*, *Br. J. Pharmacol.*, 1999, **126**: 485-493; Read *et al.*, *Cephalgia*, 1997, **17**: 826-832) or by anti-hyperalgesic properties (Bingham *et al.*, 1999 *Cephalgia* in press IHS meeting abstract).

#### Irritable Bowel Syndrome as associated with Intestinal Pain

Male Wistar rats (180-450g), housed individually and fasted overnight, were used. Under halothane (4-5%) anaesthesia, a 6-7 cm long latex balloon was inserted intra-anally to a position approximately 1 cm beyond the ano-rectal verge. The cannula from the balloon was taped to the tail to prevent its expulsion and was connected to a pressure transducer linked to a pen recorder via a 3-way tap and a peristaltic pump. Throughout the procedure the singly-housed animals were allowed unrestricted movement. After full recovery from anaesthesia, as determined by recovery of righting reflex and stable gait, a ramp inflation of the colo-rectal

5 balloon was performed until the visceromotor threshold was observed as contraction of the  
abdominal muscles. At this point the stimulus was immediately removed by opening the 3-way  
tap to air. The threshold distension pressure was noted from the pen recorder. This inflation  
10 procedure was repeated at 5 min intervals for up to 75 min, during which time constant responses  
5 are evoked with thresholds of between 10-40 mmHg. Orexin receptor antagonists, or saline were  
dosed sc after 3 stable responses to colo-rectal distension were achieved and within 45 min of  
dosing with any substance which might evoke intestinal allodynia. Further distensions were  
performed at 5 min intervals over 30 min. If, at any time, the distension pressure reached 100  
15 mmHg without observation of the visceromotor reflex then that inflation procedure was  
10 terminated and the stimulus removed. Changes in the distension pressure required to evoke the  
visceromotor reflex were compared with the mean of the three pre-dose stable recordings which  
was taken to be 100%.

**Claims**

5

10

15

20

25

30

35

40

45

50

55

5

Claims

10

1. The use of an orexin receptor antagonist in the manufacture of a medicament for the treatment and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events, the treatment and/or prophylaxis of nausea and vomiting, or the treatment and/or prophylaxis of irritable bowel syndrome or other conditions associated with visceral pain.

15

2. The use according to claim 1 wherein the medicament is for the treatment and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events.

20

3. The use according to claim 2 wherein the medicament is for the treatment and/or prophylaxis of cerebral ischaemia after cardiac arrest, stroke and multi-infarct dementia, cerebral ischaemic events resulting from surgery and/or during childbirth, or a disorder resulting from traumatic head injury and subsequent haemorrhaging.

25

4. The use according to claim 2 wherein the medicament is for the treatment and/or prophylaxis of stroke or sub-arachnoid haemorrhage.

5. The use according to claim 1 wherein the medicament is for the treatment and/or prophylaxis of nausea and vomiting.

30

6. The use according to claim 5 wherein the medicament is for the treatment and/or prophylaxis of nausea and vomiting associated with cancer therapy or post-operative or motion sickness.

35

7. The use according to claim 1 wherein the medicament is for the treatment and/or prophylaxis of irritable bowel syndrome.

8. The use according to claim 1 wherein the medicament is for the treatment of a condition associated with visceral pain.

40

9. The use according to claim 8 wherein the condition associated with visceral pain is migraine, angina or urge type incontinence.

45

10. The use according to any one of the preceding claims wherein the orexin receptor antagonist is an orexin-1 receptor antagonist.

50

55



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>A61K 31/00, A61K 31/17,</b> <b>A61K 38/00, A61P 1/00,</b> <b>A61P 1/08, A61P 9/10,</b> <b>A61P 13/10, A61P 25/06,</b> <b>A61P 25/28</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/47284</b> <b>(43) International Publication Date:</b> 17 August 2000 (17.08.2000)
--	-----------	---

**(21) International Application Number:** PCT/EP00/01147**(22) International Filing Date:** 10 February 2000 (10.02.2000)**(30) Priority Data:**

9903265.8	12 February 1999 (12.02.1999)	GB
9903278.1	12 February 1999 (12.02.1999)	GB
9903282.3	12 February 1999 (12.02.1999)	GB
9903284.9	12 February 1999 (12.02.1999)	GB
9906061.8	17 March 1999 (17.03.1999)	GB

**(60) Parent Application or Grant**

SMITHKLINE BEECHAM P.L.C. [/]; O. IRVING, Elaine,  
Alison [/]; O. SANGER, Gareth, John [/]; O. IRVING, Elaine,  
Alison [/]; O. SANGER, Gareth, John [/]; O. BLAKEY,  
Alison, Jane ; O.

**Published****(54) Title: NOVEL USE OF OREXIN RECEPTOR ANTAGONISTS****(54) Titre: NOUVELLE UTILISATION DES ANTAGONISTES DES RECEPTEURS DE L'OREXINE****(57) Abstract**

The use of orexin receptor antagonists as neuroprotectants, and in the treatment of nausea and vomiting, irritable bowel syndrome and other conditions associated with visceral pain.

**(57) Abrégé**

L'invention concerne l'utilisation des antagonistes des récepteurs de l'orexine comme neuroprotecteurs, ainsi que dans le cadre du traitement des nausées et vomissements, des colopathies fonctionnelles et autres pathologies associées à une douleur viscérale.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 August 2000 (17.08.2000)

PCT

(10) International Publication Number  
**WO 00/47284 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 31/00**,  
38/00, A61P 1/08, 9/10, 13/10, 25/28, 25/06, 1/00 // A61K  
31/17

(74) Agent: **BLAKEY, Alison, Jane**; SmithKline Beecham  
Corporate Intellectual Property, Two New Horizons Court,  
Brentford, Middlesex TW8 9EP (GB).

(21) International Application Number: **PCT/EP00/01147**

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,  
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,  
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,  
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,  
UG, US, UZ, VN, YU, ZA, ZW.

(22) International Filing Date: 10 February 2000 (10.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
9903282.3 12 February 1999 (12.02.1999) GB  
9903278.1 12 February 1999 (12.02.1999) GB  
9903284.9 12 February 1999 (12.02.1999) GB  
9903265.8 12 February 1999 (12.02.1999) GB  
9906061.8 17 March 1999 (17.03.1999) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW). Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE). OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):  
**SMITHKLINE BEECHAM P.L.C.** [GB/GB]; New  
Horizons Court, Brentford, Middlesex TW8 9EP (GB).

**Published:**

- *With international search report.*
- *Before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments.*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **IRVING, Elaine**,  
Alison [GB/GB]; SmithKline Beecham Pharmaceuticals,  
New Frontiers Science Park South, Third Avenue, Har-  
low, Essex CM19 5AW (GB). **SANGER, Gareth, John**  
[GB/GB]; SmithKline Beecham Pharmaceuticals, New  
Frontiers Science Park South, Third Avenue, Harlow,  
Essex CM19 5AW (GB).

(88) Date of publication of the international search report:  
14 December 2000

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

WO 00/47284 A3

(54) Title: **NOVEL USE OF OREXIN RECEPTOR ANTAGONISTS**

(57) Abstract: The use of orexin receptor antagonists as neuroprotectants, and in the treatment of nausea and vomiting, irritable bowel syndrome and other conditions associated with visceral pain.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/01147

A. CLASSIFICATION OF SUBJECT MATTER		
IPC 7	A61K31/00 A61P25/28	A61K38/00 A61P25/06
A61P1/08 A61P1/00	A61P9/10 //A61K31/17	A61P13/10
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC 7 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
EPO-Internal		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 849 361 A (SMITHKLINE BEECHAM CORPORATION) 24 June 1998 (1998-06-24) cited in the application page 3, line 43 - line 53 page 9, line 56 - line 58 page 10, line 1 - line 10 see page 11 lines 14,15,36-58 page 12, line 1 - line 23	1,8-10
A	---	2,3,7
X	EP 0 875 566 A (SMITHKLINE BEECHAM CORPORATION) 4 November 1998 (1998-11-04) cited in the application page 12, line 26 - line 49 page 13, line 36 - line 41 page 14, line 12 - line 33 ---	1,8-10
-/-		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "B" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
28 September 2000		06/10/2000
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patendaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Gac, G

Form PCT/ISA/210 (second sheet) (July 1992)



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 00/01147

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 875 565 A (SMITHKLINE BEECHAM CORPORATION) 4 November 1998 (1998-11-04) cited in the application page 10, line 22 - line 45 page 11, line 32 - line 37 page 12, line 10 - line 29 ---	1,8-10
X	EP 0 893 498 A (SMITHKLINE BEECHAM CORPORATION) 27 January 1999 (1999-01-27) cited in the application page 12, line 54 - line 58 page 13, line 1 - line 3 ---	1,8,9
X	WO 96 34877 A (SOPPET ET AL.) 7 November 1996 (1996-11-07) cited in the application page 1, paragraph 1 page 6, paragraph 3 page 24, paragraphs 4,5 page 26, paragraphs 2-6 ---	1,2,10
P,X	WO 99 09024 A (SMITHKLINE BEECHAM CORPORATION) 25 February 1999 (1999-02-25) cited in the application see page 2 lines 1,3,6,17-31 page 3, line 20 - line 22 page 12, line 6 - line 32 & PCT/GB 9802437 ---	1-4,8-10
P,X	WO 99 58533 A (JOHNS ET AL.) 18 November 1999 (1999-11-18) cited in the application page 2, line 26 - line 34 page 9, line 7 - line 34 ---	1,8-10
A	& PCT/EP 9903100 ---	2-4
P,X	WO 99 30670 A (SMITHKLINE BEECHAM CORPORATION) 24 June 1999 (1999-06-24) page 21, line 7 - line 33 page 26, line 31 ---	1,8-10
A	---	2,3
E,L	WO 00 27845 A (MERCK & CO. INC.) 18 May 2000 (2000-05-18) page 34, line 1 - line 9 ---	1-7,9
	---	

-/--

# INTERNATIONAL SEARCH REPORT

Inter:      nial Application No  
PCT/EP 00/01147

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	<p>KIRCHGESSNER A L (REPRINT) ET AL: "Orexin synthesis and response in the gut" NEURON, (DEC 1999) VOL. 24, NO. 4, PP. 941-951. PUBLISHER: CELL PRESS, 1050 MASSACHUSETTES AVE, CIRCULATION DEPT, CAMBRIDGE, MA 02138. ISSN: 0896-6273., XP000937996</p> <p>SUNY HLTH SCI CTR, DEPT PHYSIOL &amp; PHARMACOL, BROOKLYN, NY 11203 (Reprint);COLUMBIA UNIV COLL PHYS &amp; SURG, DEPT ANAT &amp; CELL BIOL, NEW YORK, NY 10032</p> <p>page 945, right-hand column page 948, right-hand column -----</p>	1,5,7

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-4,10

Use of an orexin antagonist in the prophylaxis or treatment of neuronal degeneration (possibly resulting from events of claim 3)

2. Claims: 5,6,10

Use of an orexin antagonist in the prophylaxis or treatment of nausea and vomiting

3. Claims: 7,10

Use of an orexin antagonist in the prophylaxis or treatment of irritable bowel syndrome

4. Claims: 8-10

Use of an orexin antagonist in the prophylaxis or treatment of migraine

5. Claims: 8-10

Use of an orexin antagonist in the prophylaxis or treatment of angina

6. Claims: 8-10

Use of an orexin antagonist in the prophylaxis or treatment of urge-type incontinence

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 1-10 relate to a product/compound defined by reference to a desirable characteristic or property, namely "an orexin (-1) receptor antagonist" and claims 1,8 to disorders only defined by a parallel/common symptom, namely "conditions associated with visceral pain".

The claims cover all products/compounds and disorders having these characteristics or properties, whereas the application provides no single support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for such products/compounds and only limited support for specific disorders.

In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely on the general concept of using orexin antagonists (without any regard to particular compounds) in the treatment or prophylaxis of neuronal degeneration (see claims 1-4), nausea and vomiting (claims 5-6), irritable bowel syndrome, (pain in general), migraine, angina, urge-type incontinence.

Claims searched completely: none

claims searched incompletely: claims 1-10

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/EP 00/01147

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 849361 A	24-06-1998	CA 2218452 A JP 10229887 A US 6001963 A	07-06-1998 02-09-1998 14-12-1999
EP 875566 A	04-11-1998	US 6020157 A DE 875566 T JP 10327888 A JP 2000060578 A	01-02-2000 04-05-2000 15-12-1998 29-02-2000
EP 875565 A	04-11-1998	US 5935814 A JP 10327889 A JP 2000106888 A	10-08-1999 15-12-1998 18-04-2000
EP 893498 A	27-01-1999	JP 11178588 A	06-07-1999
WO 9634877 A	07-11-1996	CA 2220036 A AU 715286 B AU 2470795 A EP 0828751 A JP 11505110 T	07-11-1996 20-01-2000 21-11-1996 18-03-1998 18-05-1999
WO 9909024 A	25-02-1999	AU 8741198 A EP 1003737 A	08-03-1999 31-05-2000
WO 9958533 A	18-11-1999	AU 4037799 A	29-11-1999
WO 9930670 A	24-06-1999	EP 1037655 A	27-09-2000
WO 0027845 A	18-05-2000	AU 1473200 A	29-05-2000